

Conformable Fourier Transform on Time Scales

S. G. Georgiev, T. G. Thange and S. M. Chhatraband

Abstract. In this paper, we formulate the conformable Fourier transform on time scales, drawing motivation from the structure of the conformable bilateral Laplace transform. Some of the elementary properties are proved, including shifting, transform of derivative, conjugation, transform of Hilger delta function, and transform of integral.

Key Words: Time Scale, Conformable Laplace Transform, Conformable Bilateral Laplace Transform, Conformable Fourier Transform, Generalized Exponential Function

Mathematics Subject Classification 2020: 26E70, 34A08, 34N05, 44A10

1 Introduction

Since Hilger's pioneering work [14], the study of time scales and measure chains has drawn considerable attention. Significant progress has been made by researchers, particularly in exploring dynamic equations on time scales, which unify concepts from both differential and difference equations (see, for example, [4]).

In the study of dynamic equations on time scales, the development of the integral transform method is a key focus for solving initial value problems. Thus far, various integral transforms have been generalized on time scales; we refer to [6, 9, 11, 12, 17–24] among the others.

The classical Fourier transform is a mathematical technique that converts a signal from the time domain to the frequency domain, revealing the different frequency components present and their magnitudes. During the initial development of time scale theory, Hilger [15] began exploring Fourier analysis on time scales. The Fourier transform defined by Hilger combines various forms of Fourier analysis into a unified framework. It offers a closed-form expression representing both the Fourier integral and Fourier series, through

a simple integral defined on a time scale. Later, he explored Fourier transforms on specific subclasses of time scales that also have a group structure, as discussed in [16]. The concept of discrete Fourier transform was introduced in [7], extending Fourier analysis to discrete time scales. Furthermore, a “generalized Fourier transform” with kernel as a classical exponential function is defined on a time scale.

Researchers have extended the conformable fractional calculus to arbitrary time scales by applying the principles of classical fractional calculus (see, for example, [1–3, 26]). In [25], we introduced conformable Laplace transform on time scales as follows:

$$\mathfrak{L}_\alpha\{f(t)\}(z) = \int_{t_0}^{\infty} \mathbb{E}_{\ominus_\alpha}(\sigma(t), t_0) f(t) \Delta^\alpha t.$$

Subsequently, in [13], the conformable bilateral Laplace transform was defined on time scales, where the region of integration is the entire time scales, as follows:

$$\mathcal{L}^b(f)(z, s) = \int_{-\infty}^{\infty} \mathbb{E}_{\ominus_\alpha z}^\sigma(t, s) f(t) \Delta^\alpha t. \quad (1)$$

Motivated by the bilateral Laplace transform given by equation (1), in the present work, we develop conformable Fourier transform on time scales.

This paper is organized as follows. The foundational concepts and notations of conformable fractional calculus are presented in Section 2. In Section 3, we introduce the conformable Fourier transform on time scales. Some of its basic properties, including transform of conformable fractional derivative, and conformable α -fractional integral, are given. At last, in Section 4, we solve a conformable dynamic equation using conformable Fourier transform.

2 Preliminaries

Assuming a foundational understanding of time scale calculus as elaborated in [4, 5, 8, 10, 26], we present the essential definitions and theorems for our discussion. Throughout this paper, we consider a time scale \mathbb{T} that is unbounded both above and below. For $t \in \mathbb{T}$, the forward jump operator $\sigma : \mathbb{T} \rightarrow \mathbb{T}$ is defined as

$$\sigma(t) = \inf\{\tau \in \mathbb{T} : \tau > t\},$$

and the forward graininess function $\mu : \mathbb{T} \rightarrow [0, \infty)$ is defined as

$$\mu(t) = \sigma(t) - t.$$

For a given time scale \mathbb{T} , the non-maximal set \mathbb{T}^κ is given by

$$\mathbb{T}^\kappa = \begin{cases} \mathbb{T} \setminus \rho(\sup \mathbb{T}, \sup \mathbb{T}] & \text{for } \sup \mathbb{T} < \infty, \\ \mathbb{T} & \text{for } \sup \mathbb{T} = \infty. \end{cases}$$

Definition 1 A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is called regressive if $1 + \mu(t)f(t) \neq 0$ for all $t \in \mathbb{T}^\kappa$, and a positively regressive if $1 + \mu(t)f(t) > 0$ for all $t \in \mathbb{T}^\kappa$. The set of regressive and positively regressive functions are denoted as \mathcal{R} and \mathcal{R}^+ , respectively.

Let $h > 0$. The Hilger complex numbers and the Hilger real axis are defined as

$$\mathbb{C}_h = \mathbb{C} \setminus \left\{ \frac{-1}{h} \right\} \quad \text{and} \quad \mathbb{R}_h = (\mathbb{C}_h \cap \mathbb{R}) \setminus \left\{ \frac{-1}{h} \right\},$$

respectively. If $h = 0$, we have $\mathbb{C}_0 := \mathbb{C}$ and $\mathbb{R}_0 = \mathbb{R}$. Further, \mathbb{Z}_h is a strip

$$\mathbb{Z}_h := \left\{ z \in \mathbb{C} : \frac{-\pi}{h} < \text{Im}(z) \leq \frac{\pi}{h} \right\},$$

and for $h = 0$, we have $\mathbb{Z}_0 = \mathbb{C}$.

For $z \in \mathbb{C}_h$, the Hilger real part of z is given by

$$\text{Re}_h(z) := \frac{|hz + 1| - 1}{h}.$$

For $k > 0$, the cylindrical transformation $\xi_k : \mathbb{C}_k \rightarrow \mathbb{Z}_k$ is defined as

$$\xi_k(z) = \frac{1}{k} \log(1 + zk),$$

where \log is the principal logarithm function.

Definition 2 Let $g : \mathbb{T} \rightarrow \mathbb{R}$, $t \in \mathbb{T}^\kappa$, and $\alpha \in (0, 1]$. Suppose there is a number $g^{(\alpha)}(t)$ such that for $\epsilon > 0$, there exists a neighbourhood $U \subset \mathbb{T}$ of t such that

$$|(g(\sigma(t)) - g(s))|t|^{1-\alpha} - g^{(\alpha)}(t)(\sigma(t) - s)| \leq \epsilon|\sigma(t) - s|$$

for all $s \in U$. Then $g^{(\alpha)}(t)$ is called the conformable fractional derivative of g of order α at t .

If g is a regulated function, its conformable α -fractional integral is given by

$$\int g(t) \Delta^\alpha t = \int g(t) |t|^{\alpha-1} \Delta t.$$

Definition 3 A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is said to be *rd-continuous* if f is continuous at right-dense points in \mathbb{T} and has a finite limit at the left-dense points in \mathbb{T} . We denote the set of all rd-continuous functions by $\mathcal{C}_{\text{rd}}(\mathbb{T}, \mathbb{R})$.

Definition 4 A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is said to be ‘ α -regressive’ if

$$1 + \mu(t)f(t)|t|^{\alpha-1} \neq 0 \quad \text{for all } t \in \mathbb{T}^\kappa,$$

and is said to be ‘ α -positively regressive’ provided

$$1 + \mu(t)f(t)|t|^{\alpha-1} > 0, \quad \text{for all } t \in \mathbb{T}^\kappa.$$

We denote the set of all α -regressive and rd-continuous (α -positively regressive and rd-continuous) functions by $\mathcal{R}^\alpha(\mathcal{R}^{\alpha+})$.

For a set \mathcal{R}^α , ‘ α -circle plus’ addition \oplus_α is a binary operation defined as

$$(f_1 \oplus_\alpha f_2)(t) = f_1(t) + f_2(t) + \mu(t)f_1(t)f_2(t)|t|^\alpha \quad \text{for all } t \in \mathbb{T}^\kappa.$$

The set \mathcal{R}^α forms an abelian group under \oplus_α . For $f \in \mathcal{R}^\alpha$, the inverse of f is given as

$$\ominus_\alpha f(t) = \frac{-f(t)}{1 + \mu(t)f(t)|t|^\alpha}, \quad t \in \mathbb{T}^\kappa.$$

Further, \mathcal{R}^α , ‘ α -circle minus’ subtraction \ominus_α is defined as

$$(f_1 \ominus_\alpha f_2)(t) = \frac{f_1(t) - f_2(t)}{1 + \mu(t)f_2(t)|t|^{\alpha-1}}, \quad t \in \mathbb{T}^\kappa.$$

For $f \in \mathcal{R}^\alpha$, the generalized exponential function is defined by

$$\mathbb{E}_f(t, s) = \exp \left(\int_s^t \xi_{\mu(\tau)}(f(\tau)|\tau|^{\alpha-1}) \Delta\tau \right) \quad \text{for all } t, s \in \mathbb{T}. \quad (2)$$

Applying the concept of the cylindrical transformation [4, Definition 2.21], equation (2) can be written as

$$\mathbb{E}_f(t, s) = \exp \left(\int_s^t \frac{1}{\mu(\tau)} \text{Log}(1 + \mu(\tau)f(\tau)|\tau|^{\alpha-1}) \Delta\tau \right), \quad t, s \in \mathbb{T}.$$

Some important properties of the generalized exponential function are given in the following theorem.

Theorem 1 If $g_1, g_2 \in \mathcal{R}^\alpha$, then for all $s, t, r \in \mathbb{T}$,

$$(1) \quad \mathbb{E}_0(t, s) = 1 \text{ and } \mathbb{E}_{g_1}(t, t) = 1;$$

$$(2) \quad \mathbb{E}_{g_1}(t, s)\mathbb{E}_{g_1}(s, r) = \mathbb{E}_{g_1}(t, r);$$

$$(3) \quad \mathbb{E}_{g_1}(t, s) = \frac{1}{\mathbb{E}_{g_1}(s, t)} = \mathbb{E}_{\ominus_\alpha g_1}(s, t);$$

$$(4) \quad \mathbb{E}_{g_1}(t, s)\mathbb{E}_{g_2}(t, s) = \mathbb{E}_{g_1 \oplus_\alpha g_2}(t, s);$$

$$(5) \quad \frac{\mathbb{E}_{g_1}(t, s)}{\mathbb{E}_{g_2}(t, s)} = \mathbb{E}_{g_1 \ominus_\alpha g_2}(t, s);$$

$$(6) \quad \mathbb{E}_{g_1}(\sigma(t), s) = (1 + \mu(t)g_1(t)|t|^{\alpha-1})\mathbb{E}_{g_1}(t, s);$$

$$(7) \quad \mathbb{E}_{\ominus_\alpha g_1}(\sigma(t), s) = \frac{\mathbb{E}_{\ominus_\alpha g_1}(t, s)}{1 + \mu(t)g_1(t)|t|^{\alpha-1}};$$

$$(8) \quad \mathbb{E}_{g_1}^\Delta(t, s) = g_1(t)\mathbb{E}_{g_1}(t, s)|t|^{\alpha-1};$$

$$(9) \quad \mathbb{E}_{g_1}^{(\alpha)}(t, s) = g_1(t)\mathbb{E}_{g_1}(t, s).$$

Definition 5 Let $s \in \mathbb{T}$, $\beta \in \mathcal{R}^{\alpha+}([s, \infty))$, $\gamma \in \mathcal{R}^{\alpha+}((-\infty, s])$. We say that (s, β, γ) is an admissible triple if

$$\mathbb{C}_{s, \beta, \gamma} = \left\{ z \in \mathbb{C} : \operatorname{Re}_{\mu^*(s)}(z) < \gamma, \operatorname{Re}_{\mu_*(s)}(z) > \beta, 1 + \bar{\mu}(s)\operatorname{Re}_{\bar{\mu}(s)}(z) \neq 0 \right\} \neq \emptyset.$$

Definition 6 Let $f : \mathbb{T} \rightarrow \mathbb{R}$ be regulated. Then for $s \in \mathbb{T}$, the conformable bilateral Laplace transform of f is defined by

$$\mathcal{L}^b(f)(z, s) = \int_{-\infty}^{\infty} f(t)\mathbb{E}_{\ominus_\alpha z}^\sigma(t, s)\Delta^\alpha t$$

for $z \in \mathbb{C}$ such that $1 + \mu(t)z|t|^{\alpha-1} \neq 0$ for any $t \in \mathbb{T}^\kappa$ and the improper integral exists.

Next theorems give the conditions for the absolute and uniform convergence for the conformable bilateral Laplace transform.

Theorem 2 [13] Let (s, β, γ) is an admissible triple and $f \in \mathcal{C}_{rd}(\mathbb{T})$ is of conformable double exponential order (β, γ) . Then $\mathcal{L}^b(f)(\cdot, s)$ exists on $\mathbb{C}_{s, \beta, \gamma}$ and converges absolutely.

Theorem 3 [13] Let (s, β, γ) is an admissible triple and $f \in \mathcal{C}_{rd}(\mathbb{T})$ is of conformable double exponential order (β, γ) . Then $\mathcal{L}^b(f)(\cdot, s)$ converges uniformly in $\mathbb{C}_{s, \beta, \gamma}$.

3 The conformable Fourier transform

Suppose that \mathbb{T} is a time scale with forward jump operator σ and delta differentiation operator Δ . Also, let $\inf \mathbb{T} = -\infty$, $\sup \mathbb{T} = \infty$ and $s \in \mathbb{T}$. Denote

$$\mu_*(s) = \inf_{t \in [s, \infty)} \mu(t), \quad \mu^*(s) = \sup_{t \in (-\infty, s]} \mu(t), \quad \bar{\mu}(s) = \inf_{t \in (-\infty, s]} \mu(t),$$

and for $x \in \mathbb{R}$, put

$$\bar{\bar{\mu}}(s) = \begin{cases} \mu^*(s) & \text{if } \operatorname{Re}_{\bar{\mu}(s)}(x) \leq 0, \\ \bar{\mu}(s) & \text{if } \operatorname{Re}_{\bar{\mu}(s)}(x) > 0. \end{cases}$$

Definition 7 Suppose that $f : \mathbb{T} \rightarrow \mathbb{R}$ is regulated. Then the Fourier transform of the function f is defined by

$$\mathcal{F}(f)(x, s) = \int_{-\infty}^{\infty} f(t) \mathbb{E}_{\ominus_{\alpha} ix}^{\sigma}(t, s) \Delta^{\alpha} t$$

for $x \in \mathbb{R}$ such that $1 + ix\mu(t)|t|^{\alpha-1} \neq 0$ for any $t \in \mathbb{T}^{\kappa}$ and the improper integral exists.

Definition 8 Let $\beta \in \mathcal{R}^+([s, \infty))$, $\gamma \in \mathcal{R}^+((-\infty, s])$. We say that (s, β, γ) is a real admissible triple if

$$R_{s, \beta, \gamma} = \left\{ x \in \mathbb{R} : \operatorname{Re}_{\mu^*(s)}(ix) < \gamma, \operatorname{Re}_{\mu_*(s)}(ix) > \beta, \right. \\ \left. 1 + \bar{\bar{\mu}}(s) \operatorname{Re}_{\bar{\mu}(s)}(ix) \neq 0 \right\} \neq \emptyset.$$

If $f \in \mathcal{C}_{rd}(\mathbb{T})$, then triple (s, β, γ) is a real admissible triple and f is of double exponential order (β, γ) . By Theorem 2 and Theorem 3, it follows that $\mathcal{F}(f)(\cdot, s)$ exists on $R_{s, \beta, \gamma}$ and converges absolutely and uniformly there. Below we present some of the properties of the Fourier transform.

Theorem 4 Let $f, g : \mathbb{T} \rightarrow \mathbb{R}$, $\lambda_1, \lambda_2 \in \mathbb{C}$. Then

$$\mathcal{F}(\lambda_1 f + \lambda_2 g)(x, s) = \lambda_1 \mathcal{F}(f)(x, s) + \lambda_2 \mathcal{F}(g)(x, s)$$

for those $x \in \mathbb{R}$ for which $1 + ix\mu(t)|t|^{\alpha-1} \neq 0$, $t \in \mathbb{T}^{\kappa}$, and the corresponding integrals exist.

Proof. Indeed,

$$\begin{aligned}
\mathcal{F}(\lambda_1 f + \lambda_2 g)(x, s) &= \int_{-\infty}^{\infty} (\lambda_1 f + \lambda_2 g)(t) \mathbb{E}_{\ominus_{\alpha} ix}^{\sigma}(t, s) \Delta^{\alpha} t \\
&= \lambda_1 \int_{-\infty}^{\infty} f(t) \mathbb{E}_{\ominus_{\alpha} ix}^{\sigma}(t, s) \Delta^{\alpha} t + \lambda_2 \int_{-\infty}^{\infty} g(t) \mathbb{E}_{\ominus_{\alpha} ix}^{\sigma}(t, s) \Delta^{\alpha} t \\
&= \lambda_1 \mathcal{F}(f)(x, s) + \lambda_2 \mathcal{F}(g)(x, s).
\end{aligned}$$

□

Theorem 5 Let $f : \mathbb{T} \rightarrow \mathbb{R}$. Then

$$\mathcal{F}(\mathbb{E}^{\sigma} y(\cdot, s) f(\cdot))(x, s) = \mathcal{F}(f)(z, s),$$

where $z = x + iy/1 + \mu|t|^{\alpha-1}$, and $x, y \in \mathbb{R}$ are such that $1 + \mu(t)|t|^{\alpha-1}y \neq 0$ for any $t \in \mathbb{T}^{\kappa}$, and the corresponding integrals exist.

Proof. Note that

$$iz = i \frac{x + iy}{1 + \mu|t|^{\alpha-1}y} = - \frac{x + iy}{i(1 + \mu|t|^{\alpha-1}y)}$$

and

$$\begin{aligned}
\ominus_{\alpha} iz &= \frac{-iz}{1 + i\mu|t|^{\alpha-1}z} = \frac{\frac{x+iy}{i(1+\mu|t|^{\alpha-1}y)}}{1 - \mu|t|^{\alpha-1} \frac{x+iy}{i(1+\mu|t|^{\alpha-1}y)}} \\
&= \frac{x + iy}{i + i\mu|t|^{\alpha-1}y - \mu|t|^{\alpha-1}x - i\mu|t|^{\alpha-1}y} \\
&= \frac{x + iy}{i - \mu|t|^{\alpha-1}x} = \frac{i(x + iy)}{-1 - i\mu|t|^{\alpha-1}x} \\
&= \frac{y - ix}{1 + i\mu|t|^{\alpha-1}x} = y \ominus_{\alpha} ix.
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathcal{F}(\mathbb{E}^{\sigma} y(\cdot, s) f(\cdot))(x, s) &= \int_{-\infty}^{\infty} \mathbb{E}_{\ominus_{\alpha} ix}^{\sigma}(t, s) \mathbb{E} y^{\sigma}(t, s) f(t) \Delta^{\alpha} t \\
&= \int_{-\infty}^{\infty} \mathbb{E}_{y \ominus_{\alpha} ix}^{\sigma}(t, s) f(t) \Delta^{\alpha} t \\
&= \mathcal{F}(f)(z, s).
\end{aligned}$$

□

Theorem 6 *Let $f : \mathbb{T} \rightarrow \mathbb{R}$. For any $k \in \mathbb{N}$, we have*

$$\mathcal{F}\left(f^{(\alpha^k)}\right)(x, s) = (ix)^k \mathcal{F}(f)(x, s)$$

for those $x \in \mathbb{R}$ for which $1 + ix\mu(t)|t|^{\alpha-1} \neq 0$, $t \in \mathbb{T}^\kappa$, the corresponding integrals exist, and

$$\lim_{t \rightarrow \pm\infty} f^{(\alpha^l)}(t) \mathbb{E}_{\ominus_\alpha ix}(t, s) = 0, \quad l \in \{0, \dots, k-1\}.$$

Proof. We will use the principle of mathematical induction. For $k = 1$, we have

$$\begin{aligned} \mathcal{F}\left(f^{(\alpha)}\right)(x, s) &= \int_{-\infty}^{\infty} f^{(\alpha)}(t) \mathbb{E}_{\ominus_\alpha ix}^\sigma(t, s) \Delta^\alpha t \\ &= \lim_{t \rightarrow \infty} f(t) \mathbb{E}_{\ominus_\alpha ix}(t, s) - \lim_{t \rightarrow -\infty} f(t) \mathbb{E}_{\ominus_\alpha ix} \\ &\quad - \int_{-\infty}^{\infty} (\ominus_\alpha ix) f(t) \mathbb{E}_{\ominus_\alpha ix}(t, s) \Delta^\alpha t \\ &= ix \int_{-\infty}^{\infty} f(t) \mathbb{E}_{\ominus_\alpha ix}^\sigma(t, s) \Delta^\alpha t \\ &= ix \mathcal{F}(f)(x, s). \end{aligned}$$

Further, if

$$\mathcal{F}\left(f^{(\alpha^k)}\right)(x, s) = (ix)^k \mathcal{F}(f)(x, s)$$

for some $k \in \mathbb{N}$, then

$$\mathcal{F}\left(f^{(\alpha^{k+1})}\right)(x, s) = ix \mathcal{F}\left(f^{(\alpha^k)}\right)(x, s) = (ix)^{k+1} \mathcal{F}(f)(x, s).$$

This completes the proof. \square

Theorem 7 *Let $f : \mathbb{T} \rightarrow \mathbb{R}$. Then*

$$\overline{\mathcal{F}(f)(x, s)} = \mathcal{F}(f)(-x, s)$$

for those $x \in \mathbb{R}$ for which $1 \pm ix\mu(t)|t|^{\alpha-1} \neq 0$, $t \in \mathbb{T}^\kappa$, and the corresponding integrals exist.

Proof. Let $t \in \mathbb{T}^\kappa$ and $x \in \mathbb{R}$ be such that $1 \pm ix\mu(t)|t|^{\alpha-1} \neq 0$. Then we have

$$\begin{aligned} (\ominus_\alpha(ix))(t) &= -\frac{ix}{1 + i\mu(t)|t|^{\alpha-1}x} \\ &= -\frac{ix(1 - i\mu(t)|t|^{\alpha-1}x)}{(1 + i\mu(t)|t|^{\alpha-1}x)(1 - i\mu(t)|t|^{\alpha-1}x)} \\ &= -\frac{ix + \mu(t)|t|^{\alpha-1}x^2}{1 + (\mu(t))^2|t|^{2(\alpha-1)}x^2}, \end{aligned}$$

$$\begin{aligned}
1 + \mu(t)|t|^{\alpha-1} (\Theta_\alpha(ix)) (t) &= 1 - \mu(t)|t|^{\alpha-1} \frac{ix + \mu(t)|t|^{\alpha-1}x^2}{1 + (\mu(t))^2|t|^{2\alpha-2}x^2} \\
&= \frac{1 + (\mu(t))^2|t|^{2\alpha-2}x^2 - i\mu(t)|t|^{\alpha-1}x - (\mu(t))^2|t|^{2\alpha-2}x^2}{1 + (\mu(t))^2|t|^{2\alpha-2}x^2} \\
&= \frac{1 - i\mu(t)|t|^{\alpha-1}x}{1 + (\mu(t))^2|t|^{2\alpha-2}x^2} \\
&= \frac{1}{\sqrt{1 + (\mu(t))^2|t|^{2\alpha-2}x^2}} \\
&\quad \cdot \left(\frac{1}{\sqrt{1 + (\mu(t))^2|t|^{2\alpha-2}x^2}} - i \frac{\mu(t)|t|^{\alpha-1}x}{\sqrt{1 + (\mu(t))^2|t|^{2\alpha-2}x^2}} \right)
\end{aligned}$$

and

$$\begin{aligned}
(\Theta_\alpha(i(-x))) (t) &= \frac{1}{\sqrt{1 + (\mu(t))^2|t|^{2\alpha-2}x^2}} \\
&\quad \cdot \left(\frac{1}{\sqrt{1 + (\mu(t))^2|t|^{2\alpha-2}x^2}} + i \frac{\mu(t)|t|^{\alpha-1}x}{\sqrt{1 + (\mu(t))^2|t|^{2\alpha-2}x^2}} \right).
\end{aligned}$$

Let

$$r(t) = \frac{1}{\sqrt{1 + (\mu(t))^2|t|^{2\alpha-2}x^2}}, \quad \cos \theta(t) = \frac{1}{\sqrt{1 + (\mu(t))^2|t|^{2\alpha-2}x^2}}.$$

Then

$$1 + \mu(t)|t|^{\alpha-1} (\Theta_\alpha(ix)) (t) = r(t)e^{-i\theta(t)}$$

and

$$1 + \mu(t)|t|^{\alpha-1} (\Theta_\alpha(i(-x))) (t) = r(t)e^{i\theta(t)}.$$

Also,

$$\log (1 + \mu(t)|t|^{\alpha-1} (\Theta_\alpha(ix)) (t)) = \ln(r(t)) - i(\theta + 2k\pi),$$

$$\log (1 + \mu(t)|t|^{\alpha-1} (\Theta_\alpha(i(-x))) (t)) = \ln(r(t)) + i(\theta + 2k\pi), \quad k \in \mathbb{Z}.$$

From here and from the definition of the Fourier transform, we have

$$\begin{aligned}
\mathcal{F}(f)(x, s) &= \int_{-\infty}^{\infty} f(t) \mathbb{E}_{\Theta_\alpha(ix)}^\sigma(t, s) \Delta^\alpha t \\
&= \int_{-\infty}^{\infty} e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} \log(1 + \mu(\tau)|\tau|^{\alpha-1} (\Theta_\alpha(ix))(\tau)) \Delta^\alpha \tau} f(\tau) \Delta^\alpha \tau \\
&= \int_{-\infty}^{\infty} e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} (\ln r(\tau) - i(\theta(\tau) + 2k\pi)) \Delta^\alpha \tau} f(\tau) \Delta^\alpha \tau
\end{aligned}$$

and

$$\begin{aligned}
\overline{\mathcal{F}(f)(x, s)} &= \overline{\int_{-\infty}^{\infty} e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} (\ln r(\tau) - i(\theta(\tau) + 2k\pi)) \Delta^\alpha \tau} f(t) \Delta^\alpha t} \\
&= \int_{-\infty}^{\infty} \overline{e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} (\ln r(\tau) - i(\theta(\tau) + 2k\pi)) \Delta^\alpha \tau} f(t) \Delta^\alpha t} \\
&= \int_{-\infty}^{\infty} e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} (\ln r(\tau) - i(\theta(\tau) + 2k\pi)) \Delta^\alpha \tau} \overline{f(t) \Delta^\alpha t} \\
&= \int_{-\infty}^{\infty} e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} (\ln r(\tau) - i(\theta(\tau) + 2k\pi)) \Delta^\alpha \tau} f(t) \Delta^\alpha t \\
&= \int_{-\infty}^{\infty} e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} (\ln r(\tau) - i(\theta(\tau) + 2k\pi)) \Delta^\alpha \tau} f(t) \Delta^\alpha t \\
&= \int_{-\infty}^{\infty} e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} (\ln r(\tau) + i(\theta(\tau) + 2k\pi)) \Delta^\alpha \tau} f(t) \Delta^\alpha t \\
&= \int_{-\infty}^{\infty} e^{\int_s^{\sigma(t)} \frac{1}{\mu(\tau)|\tau|^{\alpha-1}} \log(1 + \mu(\tau)|\tau|^{\alpha-1} (\ominus_\alpha(i(-x))))(\tau)) \Delta^\alpha \tau} f(\tau) \Delta^\alpha \tau \\
&= \int_{-\infty}^{\infty} f(t) \mathbb{E}_{\ominus_\alpha(i(-x))}^\sigma(t, s) \Delta^\alpha t \\
&= \mathcal{F}(f)(-x, s).
\end{aligned}$$

□

Theorem 8 *Let $f : \mathbb{T} \rightarrow \mathbb{R}$ be regulated and*

$$F(t) = \int_a^t f(\tau) \Delta^\alpha \tau, \quad t \in \mathbb{T},$$

for some fixed $a \in \mathbb{T}$. Then

$$\mathcal{F}(F)(x, s) = -\frac{i}{x} \mathcal{F}(f)(x, s)$$

for those $x \in \mathbb{R}$, $x \neq 0$, for which

$$\lim_{t \rightarrow \pm\infty} F(t) \mathbb{E}_{\ominus_\alpha ix}(t, s) = 0.$$

Proof. Let $x \in \mathbb{R}$ satisfy the conditions of the theorem. Then

$$\begin{aligned}
\mathcal{F}(F)(x, s) &= \int_{-\infty}^{\infty} F(t) \mathbb{E}_{\ominus_{\alpha} ix}^{\sigma}(t, s) \Delta^{\alpha} t \\
&= \int_{-\infty}^{\infty} F(t) (1 + \mu(t) |t|^{\alpha-1} (\ominus_{\alpha}(ix))(t)) \mathbb{E}_{\ominus_{\alpha} ix}(t, s) \Delta^{\alpha} t \\
&= \int_{-\infty}^{\infty} F(t) \frac{1}{1 + i\mu(t) |t|^{\alpha-1} x} \mathbb{E}_{\ominus_{\alpha} ix}(t, s) \Delta^{\alpha} t \\
&= -\frac{1}{ix} \int_{-\infty}^{\infty} F(t) \frac{-ix}{1 + i\mu(t) |t|^{\alpha-1} x} \mathbb{E}_{\ominus_{\alpha} ix}(t, s) \Delta^{\alpha} t \\
&= \frac{i}{x} \int_{-\infty}^{\infty} F(t) (\ominus_{\alpha} ix)(t) \mathbb{E}_{\ominus_{\alpha} ix}(t, s) \Delta^{\alpha} t \\
&= \frac{i}{x} \int_{-\infty}^{\infty} F(t) \mathbb{E}_{\ominus_{\alpha} ix}^{(\alpha)}(t, s) \Delta^{\alpha} t \\
&= \frac{i}{x} \left(\lim_{t \rightarrow \infty} F(t) \mathbb{E}_{\ominus_{\alpha} ix}(t, s) - \lim_{t \rightarrow -\infty} F(t) \mathbb{E}_{\ominus_{\alpha} ix}(t, s) \right) \\
&\quad - \frac{i}{x} \int_{-\infty}^{\infty} f(t) \mathbb{E}_{\ominus_{\alpha} ix}^{\sigma}(t, s) \Delta^{\alpha} t \\
&= -\frac{i}{x} \mathcal{F}(f)(x, s).
\end{aligned}$$

□

4 Applications

In this section, we will give some applications of the conformable Fourier transform. Consider the equation

$$u^{(\alpha^n)} + a_1 u^{(\alpha^{n-1})} + \dots + a_{n-1} u^{(\alpha)} + a_n u = f(t), \quad t \in \mathbb{T},$$

where $a_j, j \in \{1, \dots, n\}$, are given constants, and $f : \mathbb{T} \rightarrow \mathbb{R}$ is given function such that $\mathcal{F}(f)(x, s)$ exists. Applying the conformable Fourier transform of both sides of the considered equation, we get

$$\begin{aligned}
\mathcal{F}(f)(x, s) &= \mathcal{F}(u^{(\alpha^n)})(x, s) + a_1 \mathcal{F}(u^{(\alpha^{n-1})})(x, s) + \dots \\
&\quad + a_{n-1} \mathcal{F}(u^{(\alpha)})(x, s) + a_n \mathcal{F}(u)(x, s) \\
&= (ix)^n \mathcal{F}(u)(x, s) + (ix)^{n-1} a_1 \mathcal{F}(u)(x, s) + \dots \\
&\quad + i x a_{n-1} \mathcal{F}(u)(x, s) + a_n \mathcal{F}(u)(x, s) \\
&= ((ix)^n + (ix)^{n-1} a_1 + \dots + i x a_{n-1} + a_n) \mathcal{F}(u)(x, s),
\end{aligned}$$

whereupon

$$\mathcal{F}(u)(x, s) = \frac{\mathcal{F}(f)(x, s)}{(ix)^n + (ix)^{n-1}a_1 + \dots + ixa_{n-1} + a_n}.$$

Taking the inverse conformable Fourier transform for the solution u of the considered equation, we obtain

$$u(t) = \mathcal{F}^{-1}\left(\frac{\mathcal{F}(f)(x, s)}{(ix)^n + (ix)^{n-1}a_1 + \dots + ixa_{n-1} + a_n}\right)(t), \quad t \in \mathbb{T}.$$

For example let us consider the equation

$$u^{(\alpha)} - u = f(t), \quad t \in \mathbb{T},$$

where

$$f(t) = \begin{cases} E_i(t, s) & \text{for } t \geq s, \\ 0 & \text{for } t < s. \end{cases}$$

We will search a solution of this equation in the form

$$u(t) = \begin{cases} v(t) & \text{for } t \geq s \\ 0 & \text{for } t < s. \end{cases}$$

Applying the conformable Fourier transform of both sides of the considered equation, we get

$$\begin{aligned} \frac{1}{x-i} &= \mathcal{F}(f)(x, s) = \mathcal{F}(v^{(\alpha)})(x, s) - \mathcal{F}(v)(x, s) \\ &= ix\mathcal{F}(v)(x, s) - \mathcal{F}(v)(x, s) = (ix-1)\mathcal{F}(v)(x, s) \\ &= i(x+1)\mathcal{F}(v)(x, s), \end{aligned}$$

whereupon

$$\mathcal{F}(v)(x, s) = -\frac{i}{(x-i)(x+i)} = -\frac{i}{x^2+1}.$$

Now, taking the inverse conformable Fourier transform, we find

$$v(t) = -i\mathcal{F}^{-1}\left(\frac{1}{x^2+1}\right)(t) = -i \sin_{1,\alpha}(t, s), \quad t \geq s.$$

Consequently,

$$u(t) = \begin{cases} -i \sin_{1,\alpha}(t, s) & \text{for } t \geq s, \\ 0 & \text{for } t < s, \end{cases}$$

is a solution of the considered equation.

References

- [1] D. R. Anderson and S. G. Georgiev, *Conformable dynamic equations on time scales*, Chapman and Hall/CRC, 2020. <https://doi.org/10.1201/9781003057406>
- [2] N. Benkhettou, S. Hassani and D.F.M. Torres, A conformable fractional calculus on arbitrary time scales. *J. King. Saud Uni.*, **28** (2016), no. 1, pp. 93–98. <https://doi.org/10.1016/j.jksus.2015.05.003>
- [3] B. Bendouma and A. Hammoudi, A nabla conformable fractional calculus on time scales. *Electron J. Math. Anal. Appl.*, **7** (2019), no. 1, pp. 202–216. <https://doi.org/10.21608/ejmaa.2019.312753>
- [4] M. Bohner and A. Peterson, *Dynamic equations on time scales: an introduction with applications*, Birkhäuser, Boston, Mass USA, 2001.
- [5] M. Bohner, G.Sh. Guseinov and B. Karpuz, Properties of the Laplace transform on time scales with arbitrary graininess. *Integral Transforms Spec. Funct.*, **22** (2011), no. 11, pp. 785–800. <https://doi.org/10.1080/10652469.2010.548335>
- [6] T. Cuchta and S. Georgiev, Analysis of the Bilateral Laplace transform on time scales with applications. *Int. J. Dyn. Syst. Differ. Equ.*, **11** (2021), no. 3-4, pp. 255–274. <https://doi.org/10.1504/IJDSDE.2021.117356>
- [7] J.M. Davis, I.A. Gravagne and R.J. Marks II, Time scale discrete Fourier transforms. *Proceedings of the Southeastern Symposium on System Theory*, **42** (2010), pp. 102–110. <https://doi.org/10.1109/SSST.2010.5442859>
- [8] J.M. Davis, I.A. Gravagne, B.J. Jackson, R.J. Marks II and A.A. Ramos, The Laplace transform on time scales revisited. *J. Math. Anal. Appl.*, **332** (2007), no. 2, pp. 1291–1307. <https://doi.org/10.1016/j.jmaa.2006.10.089>
- [9] J.M. Davis, I.A. Gravagne and R.J. Marks II, Bilateral Laplace transform on time scales: convergence, convolution, and the characterization of stationary stochastic time series. *Circuits. Syst. Signal Process.*, **29** (2010), pp. 1141–1165. <https://doi.org/10.1007/s00034-010-9196-2>
- [10] S.G. Georgiev and K. Zennir, *Advances in fractional dynamic inequalities on time scales*, World Scientific Publishing, 2003.
- [11] S. Georgiev, *The Laplace transform on time scales with applications*, LAP LAMBERT Academic Publisher, 2022.

- [12] S.G. Georgiev and V. Darvish, The generalized Fourier convolution on time scales. *Integral Transform Spec. Funct.*, **34** (2023), no. 3, pp. 211–227. <https://doi.org/10.1080/10652469.2022.2105323>
- [13] S.G. Georgiev, S.M. Chhatraband and T.G. Thange, Conformable bilateral Laplace transform on time scales (submitted).
- [14] S. Hilger, Ein Maßkettenkalkül mit Anwendung auf Zentrumsmannigfaltigkeiten, Ph.D. Thesis, Universität Würzburg, Germany, 1988.
- [15] S. Hilger, Special functions, Laplace and Fourier transform on measure chains. *Dyn. Syst. Appl.*, **8** (1999), pp. 471–488.
- [16] S. Hilger, An application of calculus on measure chains to Fourier theory and Heisenberg’s uncertainty principle. *The J. Diff. Equations Appl.*, **8** (2002), no. 10, pp. 897–936. <https://doi.org/10.1080/1023619021000000960>
- [17] R.J. Marks II, I.A. Gravagne and J.M. Davis, A generalized Fourier transform and convolution on time scales. *J. Math. Anal. Appl.*, **340** (2008), no. 2, pp. 901–919. <https://doi.org/10.1016/j.jmaa.2007.08.056>
- [18] M.D. Ortiguera, D.F.M. Torres and J.I. Trujilla, Exponentials and Laplace transforms on nonlinear time scales. *Commun. Nonlinear Sci. Numer. Simul.*, **39** (2016), pp. 252–270. <https://doi.org/10.1016/j.cnsns.2016.03.010>
- [19] T.G. Thange and S.M. Chhatraband, Laplace-Sumudu integral transform on time scales. *South East Asian J. Math. Math. Sci.*, **19** (2023), no. 1, pp. 91–102. <https://doi.org/10.56827/seajmms.2023.1901.9>
- [20] T.G. Thange and S.M. Chhatraband, A new α -Laplace transform on time scales. *Jñānābha*, **53** (2024), no. 2, pp. 151–160. <https://doi.org/10.58250/jnanabha.2023.53218>
- [21] T.G. Thange and S.M. Chhatraband, On Nabla Shahe transform and its applications. *Journal of Fractional Calculus and Applications*, **15** (2024), no. 1, pp. 1–13. <https://doi.org/10.21608/jfca.2024.229466.1029>
- [22] T.G. Thange and S.M. Chhatraband, New general integral transform on time scales. *J. Mathe. Model.*, **12** (2024), no. 4, pp. 655–669. <https://doi.org/10.22124/jmm.2024.27193.2400>
- [23] T.G. Thange and S.M. Chhatraband, Double Shehu transform for time scales with applications. *J. Classical Anal.*, **25** (2025), no. 1, pp. 75–94. <http://dx.doi.org/10.7153/jca-2025-25-05>

- [24] T.G. Thange and S.M. Chhatraband, On n -dimensional integral transform for time scales. Palest. J. Math., **14** (2025), no. 2, pp. 777–798.
- [25] T.G. Thange, S.M. Chhatraband and S.G. Georgiev, Conformable Laplace transform on time scales (submitted).
- [26] A. Younus, K. Bukhsh, M.A. Alqudah and T. Abdeljawad, Generalized exponential function and initial value problem for conformable dynamic equations. AIMS Mathematics, **7** (2012), no. 7, pp. 12050–12076. <https://doi.org/10.3934/math.2022670>

Svetlin G. Georgiev
Department of Mathematics,
Sorbonne University,
Paris, France
 svetlinggeorgiev1@gmail.com

Tukaram G. Thange
Department of Mathematics,
Yogeshwari Mahavidyalaya,
Ambajogai (M.S.), India – 431517.
 tgthange@gmail.com

Sneha M. Chhatraband
Department of Mathematics,
Dr. Babasaheb Ambedkar Marathwada University,
Chh. Sambhajinagar (M.S.), India – 431004.
School of Humanities and Engineering Sciences
MIT Academy of Engineering,
Alandi (D), Pune (M.S.), India – 412105.
 sneha.chhatrabandsneha@mitaoe.ac.in

Please, cite to this paper as published in
 Armen. J. Math., V. **17**, N. 9(2025), pp. 1–15
<https://doi.org/10.52737/18291163-2025.17.9-1-15>